



Protein Interactions *in vivo*


In one direction



$$mv^2/2 = kT/2$$

$$v = (kT/m)^{1/2}$$

Protein Interactions *in vivo*





Σε μια χημική αντίδραση που είναι αρκετά γρήγορη

Ο ρυθμός ενός ενζύμου εξαρτάται από το πόσο γρήγορα το υπόστρωμα μπορεί να διαχυθεί στην ενεργό θέση και το πόσο γρήγορα το προϊόν μπορεί να διαχυθεί έξω.

Αυτός ο ρυθμός είναι ανάλογος με τον ρυθμό διάχυσης-σύγκρουσης, ο οποίος με τη σειρά του εξαρτάται από το μέγεθος των μορίων και το ιξώδες του μέσου.

collision rate



k_{cat}/K_M is the catalytic efficiency. It is used to rank enzymes. A big k_{cat}/K_M means that an enzyme binds tightly to a substrate (small K_M), with a fast reaction of the ES complex.

k_{cat}/K_M is an apparent second order rate constant

$$v = k_{cat}/K_M [E]_0 [S]$$

k_{cat} είναι η καταλυτική σταθερά για τη μετατροπή του υποστρώματος σε προϊόν

K_M είναι η σταθερά Michaelis

- k_{cat}/K_M is the specificity constant. It is used to distinguish and describe various substrates.

TABLE 4.1 Values of $k_{\text{cat}}/K_{\text{m}}$ for some enzymes

Enzyme	Substrate	$k_{\text{cat}}/K_{\text{m}}$ ($\text{M}^{-1} \text{s}^{-1}$)
Acetylcholinesterase	Acetylcholine	1.5×10^8
Carbonic anhydrase	Carbon dioxide	8.3×10^7
Catalase	Hydrogen peroxide	4.0×10^8
Fumarase	Fumarate	1.6×10^8
Fumarase	Malate	3.6×10^7
Superoxide dismutase	Superoxide	2.8×10^9
Triosephosphate isomerase	Dihydroxyacetone phosphate	7.5×10^5
Triosephosphate isomerase	Glyceraldehyde 3-phosphate	2.4×10^8
Lysozyme	(NAG-NAM) ₃	83
Glucose isomerase	Glucose	7.4

Abbreviation: NAG-NAM, *N*-acetylglucosamine–*N*-acetylmuramic acid disaccharide.

rate of diffusion




Figure 4.2 How Proteins Work (©2012 Garland Science)

Dielectric constant

$$F = q_1 q_2 / 4\pi \epsilon r^2,$$

ϵ είναι η διηλεκτρική σταθερά


Ionic strength

$$\mu = \frac{1}{2} \sum z_i^2 C_i$$

Electrostatic screening

$$r = \left(\frac{\epsilon K T}{2 N_0 e^2 \mu} \right)^{1/2}$$

Acetylcholinesterase



Collision rates

electrostatic orientational steering





Figure 4.3 How Proteins Work (©2012 Garland Science)




Figure 4.4 How Proteins Work (©2012 Garland Science)

Cytochrome c




Figure 4.5.1 How Proteins Work (©2012 Garland Science)

Red basic
Green acidic









Figure 4.5 How Proteins Work (©2012 Garland Science)


Cytochrome c



Electrostatic steering



έναν κρίσιμο μηχανισμό μέσω του οποίου το κυτόχρωμα c (cyt c) εντοπίζει και συνδέεται αποτελεσματικά με τους φυσιολογικούς του συνεργάτες, όπως η κυτόχρωμα c αναγωγή (Σύμπλεγμα III) και η κυτόχρωμα c οξειδάση (Σύμπλεγμα IV), στον ενδομεμβρανικό χώρο των μιτοχονδρίων.







Protein pIs




Figure 4.7 How Proteins Work (©2012 Garland Science)




Figure 4.8 How Proteins Work (©2012 Garland Science)


σύμπλεγμα συνάντησης

encounter complex

(a)



(b)



(c)








Figure 4.6 How Proteins Work (©2012 Garland Science)

σύμπλεγμα συνάντησης



barnase with inhibitor barstar



inside of a cell




Figure 4.9 How Proteins Work (©2012 Garland Science)





Figure 4.10 How Proteins Work (©2012 Garland Science)



Figure 1.35 Molecular Biology of Assemblies and Machines (© Garland Science 2016)

volume exclusion



relative sizes

The collision rate thus depends most critically on the size r_A of the diffusing molecule:




Figure 4.15 How Proteins Work (©2012 Garland Science)

Find the partner

Processivity decreases the off-rate from polymeric substrates






Figure 4.16 How Proteins Work (©2012 Garland Science)





Glycogen

Table 1
Overview of differences in glycogen levels and metabolism in brain, skeletal muscle, and liver

Specialization of different tissues and cells types has led to the diversification of glycogen metabolism regulation. An overview of the main differences in glycogen content, inter- and intracellular localization, and glycogen-related regulatory enzyme expression in brain, muscle, and liver are presented.

Attributes	Brain	Skeletal muscle	Liver
Average particle size (inner diameter, nm)	10–30	10–40	110–290
Glycogen concentration (human, $\mu\text{mol/g}$ wet weight)	3–10	30–100	100–500
Estimated % of tissue weight	0.1	1–2	6–8
Estimated whole organ content (human, fed state, g)	0.5–1.5	100	400
Tissue localization	Regional variability. Gray matter > white matter	Muscle type-dependent. Type II > type I	Uniform
Cellular/subcellular localization	Cell-dependent, highest in astrocytes. Greater in areas with high synaptic density, primary branches and fine processes	Subsarcolemmal > myofibrillar	Hepatocytes, subcellular location modulated by metabolic conditions
Glycogenin isoform	GN1	GN1	GN2
Glycogen phosphorylase isoform	GPB	GPM	PGP1
Glycogen synthase isoform	GSI	GSI	GS2

Glycogen granule and the actin-rich spherical structures



Glycogen associated proteins


Primary proteins in the glycogen granule proteome and their interactions

List of main human glycosome-associated proteins as identified by name and UNIPROT identifier (ID) as well as primary interactions relevant to glycogen metabolism. Protein-protein interactions were derived from databases within UniProt (uniprot.org) (80), mainly The Biological General Repository for Interaction Datasets (BioGrid) (81) and the Protein Interaction Database and Analysis System (IntAct) (82). Abbreviations used are as follows: AMPK, 5'-AMP-activated protein kinase; EMP2A, laforin; GBE, branching enzyme; GDE, debranching enzyme; GN, glycogenin; GP, glycogen phosphorylase; GS, glycogen synthase; PhK, phosphorylase kinase; PP1, protein phosphatase; STBD1, starch-binding domain-containing protein 1; TRIM7, tripartite motif-containing protein.

Protein	Role	UniProt ID	Key glycogen-related interactions
Glycogenin (GN)	Initiation	P46976 (GN1, muscle) O15488 (GN2, liver)	GS, AMPK, GBE, GP, STBD1, PP1 (PPP1R3C, PPP1CA, PPP1R5), TRIM7
Tripartite motif-containing protein (TRIM7, GNIP)	Initiation, regulation	Q9Q029 (TRIM7)	GN
Glycogen synthase (GS)	Synthesis	P13807 (GS1, muscle) P54840 (GS2, liver)	GN, AMPK, GBE, PP1 (PPP1R3B, PPP1R3C, PPP1CA), STBD1, KAPCA, CSK21, MAPKAPK2, GSK3, PAST, laforin, MLP3B/3C, DYRK
Glycogen branching enzyme (GBE)	Synthesis	Q04446	GP, GS, GN, STBD1, GBE, VAPA
Glycogen phosphorylase (GP)	Degradation	P11217 (PYGM, muscle) P11216 (PYGB, brain) P06737 (PYGL, liver)	AMPK, PKC, GBE1, PP2, MAPKAPK2, PP1
Glycogen debranching enzyme (GDE, AGL)	Degradation	P35573	AMPK, PP1, malin, AMPK, STBD1
Malin (E3 ubiquitin-protein ligase NHLRC1)	Ubiquitin ligase	Q6VVB1	Laforin, GS, PP1, GDE, AMPK
5'-AMP-activated protein kinase (AMPK)	Kinase	P54646 (α2) Q9Y478 (β1) O43741 (β2)	Laforin, PP1 (PPP2CA, PPP2R1B), PHKG2, CAMK, GN
Laforin (EMP2A)	Carbohydrate phosphatase, ubiquitin ligase	O95278	AMPK, PP1 (PPP1R3C, PPP1R3D), GSK3B, STBD1, GS, malin,
Protein phosphatase 1 (PP1) and targeting subunits	Phosphatase, main regulatory and catalytic subunits	PP1	AMPK, laforin, GSK3B, GN, GS
		Q16821 (PPP1R3A, GM) Q86X16 (PPP1R3B, GL) Q0VCR4 (PPP1R3C, PTG) O95685 (PPP1R3D, PPP1R6) P67775 (PPP2CA) P62136 (PPP1CA) P30154 (PPP2R1B) P15735 (γ, liver, testis) Q16816 (γ, muscle) O95210	
Phosphorylase kinase (PhK)	Kinase		AMPK, PP1 (PPP1R3B), GP
Starch-binding domain-containing protein 1 (STBD1)	Cargo receptor for glycogen		GDE, GBE, PP1, malin, GN, GS, AMPK, GP, MLP3B/3C


Find the partner

glycogen synthase (GS)
glycogen-branching enzyme (GBE)




Find the partner

Cytosolic glycogen degradation




Lysosomal and Proteasome-dependent glycogen granule degradation



Plastics

Plastic	Applications	Usage Time	Degradation Time *
PET (Polyethylene terephthalate)	Bottles and other plastic containers	1–3 years	500–1000 years
HDPE (high-density polyethylene)	Pipelines, bottles	5–35 years	250–5000 years
LDPE (low-density polyethylene)	Plastic wrappers and bags	1–3 years	150 years
PVC (Polyvinyl chloride)	Pipelines and other uses in construction	35 years	>1000 years
PP (Polypropylene)	Textiles, packaging, automotive components	5–15 years	50–800 years
PHAs (Polyhydroxyalkanoates)	Bags, packaging, medical implants	-	<1 year



The general structure of PHAs



- Similar properties to conventional plastics
- Accumulated in the form of granules in microorganisms, as a type of carbon storage
 - PHB is the most studied PHA

Chain	x	R	Carbon	PHA	Abbreviation
Short	1	Hydrogen	C ₃	Poly(3-hydroxypropionate)	P(3HP)
		Methyl	C ₄	Poly(3-hydroxybutyrate)	P(3HB)
		Ethyl	C ₅	Poly(3-hydroxyvalerate)	P(3HV)
	2	Hydrogen	C ₄	Poly(4-hydroxybutirate)	P(4HB)
		Methyl	C ₅	Poly(4-hydroxyvalerate)	P(4HV)
		Hydrogen	C ₅	Poly(5-hydroxyvalerate)	P(5HV)
Medium	1	Propyl	C ₆	Poly(3-hydroxyhexanoate)	P(3HHx)
		Butyl	C ₇	Poly(3-hydroxyheptanoate)	P(3HHp)
		Pentyl	C ₈	Poly(3-hydroxyoctanoate)	P(3HO)
		Hexyl	C ₉	Poly(3-hydroxynonanoate)	P(3HN)
		Heptyl	C ₁₀	Poly(3-hydroxydecanoate)	P(3HD)
		Octyl	C ₁₁	Poly(3-hydroxyundecanoate)	P(3HUD)
		Nonyl	C ₁₂	Poly(3-hydroxydodecanoate)	P(3HDD)
		Decyl	C ₁₃	Poly(3-hydroxytridecanoate)	P(3HTD)
	2	Undecyl	C ₁₄	Poly(3-hydroxytetradecanoate)	P(3HTTD)
		Ethyl	C ₆	Poly(4-hydroxycarpoate)	P(4HC)
	3	Methyl	C ₆	Poly(5-hydroxycarpoate)	P(5HC)
		Ethyl	C ₇	Poly(4-hydroxyheptanoate)	P(4HH)
Long	1	Dodecyl	C ₁₅	Poly(3-hydroxypentadecanoate)	P(3HPD)
		Tridecyl	C ₁₆	Poly(3-hydroxyhexadecanoate)	P(3HHxD)
		Pentadecyl	C ₁₈	Poly(3-hydroxyoctadecanoate)	P(3HOD)



Find the partner




I. Kanavaki, A. Drakonaki, E.D. Geladas, A. Spyros, H. Xie, G. Tsiotis, Polyhydroxyalkanoate (PHA) Production in *Pseudomonas* sp. phDV1 Strain Grown on Phenol as Carbon Sources., *Microorganisms*. 9 (2021).

A Drakonaki, E Mathioudaki, ED Geladas, E Konsolaki, N Vitsaxakis,, N & Tsiotis, G. Production of Polyhydroxybutyrate by Genetically Modified *Pseudomonas* sp. phDV1: A Comparative Study of Utilizing Wine Industry Waste as a Carbon Source *Microorganisms* 11 (6), 1592

Carbonosomes



Negative-Stain TEM




(Jendrossek et al., 2007, Applied and Environmental Microbiology)

22


Cryo-EM

Cryo-EM Plunge Freezing




Krios G4 Cryo-TEM

Negative-Stain TEM






Cryo-EM




25


PHB granule organization




P_{xylose} *phaA* *phaB* (aceto-acetylCoA transferase and reductase)



PHB granules





Find the partner



Polyphosphate (polyP) is a linear polymer of phosphate residues linked by energy-rich phospho-anhydride bonds.

Polyphosphate granules



Polyphosphate granules









Fig. 2. TEM micrographs of thin sections from A) 10 PI, B) 20 PI and C) 40 PI, the arrows indicate the electron-rich inclusions (bar: 200 nm).



Polyphosphate granules




cellulose




cellulase

catalytic
domain



cellulose-binding
module



cellulose






Figure 4.17 How Proteins Work (©2012 Garland Science)






Figure 4.18 How Proteins Work (©2012 Garland Science)

Searching is faster in two dimensions





Figure 4.20 How Proteins Work (©2012 Garland Science)

Searching is faster in two dimensions


a

Agonist binding

G protein coupling and nucleotide exchange


Activated G protein subunits regulate effector proteins

GTP hydrolysis and inactivation of $G\alpha$ protein



GPCR-mediated G protein activation process


A



Searching is slightly faster again in one dimension

Oct-1 has two DNA-binding domains

(a)




(b)



Figure 4.22 How Proteins Work (©2012 Garland Science)

Searching is faster in smaller compartments



Searching is faster in smaller compartments






Figure 1.5.1b Molecular Biology of Assemblies and Machines (© Garland Science 2016)

Molecular landscape of *Chlamydomonas* mitochondria




Subtomogram average of the native *Chlamydomonas* respirasome

The $I_2 III_4 IV_6$ respirasome



● complex I ● complex III ● complex IV



Sticky arms






Figure 4.23 How Proteins Work (©2012 Garland Science)

tyrosine kinases receptor




tyrosine kinases receptor




SH2 domains

(a)




© 1999 GARLAND PUBLISHING INC.
A member of the Taylor & Francis Group

Structural features of recognition by SH2 domains




Structural water molecules mediating protein-peptide interactions


Lck SH2



Src SH2



BRDG1 SH2



SH3 domains





Figure 4.24 How Proteins Work (©2012 Garland Science)



©1999 GARLAND PUBLISHING INC.
A member of the Taylor & Francis Group




Figure 4.26 How Proteins Work (©2012 Garland Science)

SH3 domain







Figure 4.27 How Proteins Work (©2012 Garland Science)



deactivation and activation of kinases








Figure 4.35 How Proteins Work (©2012 Garland Science)

proline-serine-threonine-alanine-isoleucine-arginine-glutamate (PSTAIRE)

Src tyrosine kinase




Src tyrosine kinase




Active vs inactive


(a) active




(b) inactive





(A) Inactive



(B) Active



Abl kinase



Post-translational modifications of proteins




TABLE 4.3 A few covalent modifications of proteins		
Modification	Site	Comments
Phosphorylation	Ser, Thr, Tyr	Regulates activity. Regulates assembly
Acetylation	Lys	Creates part of histone code in chromatin
Methylation	Lys	Creates part of histone code in chromatin
Methylation	Arg	
Lipid attachment	Cys, C terminus	Attaches protein to membrane
SUMOylation	Lys	Role in transport, transcriptional regulation, apoptosis
Ubiquitylation	Lys	Regulates transport and degradation, plus histone readout
Limited proteolysis		Activates proteases (zymogens) in extracellular location (e.g. chymotrypsin); activates hormones (e.g. insulin)
Attachment of N-acetylglucosamine	Ser, Thr	Regulates activity in enzymes involved in glucose metabolism
Glycosylation	Asn, Ser/Thr	Eukaryotes. Recognition, membrane protein folding
Hydroxylation	Pro	Collagen: to facilitate triple helix formation. Irreversible
ADP ribosylation	Arg, Glu, Asp	As part of signaling, DNA repair and apoptosis
Sulfation	Tyr	Irreversible and probably required for activity
Carboxylation	Glu	Creates γ -carboxyglutamate (Gla), a calcium ligand

Table 4.3 How Proteins Work (©2012 Garland Science)

Phosphorylation




Figure 4.33 How Proteins Work (©2012 Garland Science)




Figure 4.34 How Proteins Work (©2012 Garland Science)

cell cycle







Figure 4.9.3 How Proteins Work (©2012 Garland Science)


cell cycle




CD kinase



CD kinase




(a)




Cyclin A - CDK2

(b)

© 1999 GARLAND PUBLISHING INC.
A member of the Taylor & Francis Group



(b)



© 1999 GARLAND PUBLISHING INC.
A member of the Taylor & Francis Group

ubiquitylation




TABLE 4.4 Dependence of half-life of cytoplasmic proteins on their N-terminal residue	
N-terminal residue	Half-life
Met, Gly, Ala, Ser, Thr, Val	> 20 h
Ile, Glu	30 min
Tyr, Gln	10 min
Pro	7 min
Leu, Phe, Asp, Lys	3 min
Arg	2 min

Table 4.4 How Proteins Work (©2012 Garland Science)






Figure 4.9.2 How Proteins Work (©2012 Garland Science)

ubiquitylation



Phosphatases









- Inducing conformational change
- Providing platform for phospho-binding proteins

Changes in Protein





- Stability
- Activity
- Interaction
- Subcellular localization

Lipidation

Lipid Structure	Effects on proteins	Effects on downstream signaling in cancer
GPI anchor 	Plasma membrane tethering Incorporation into specific membrane domains Protein-protein interaction	Increase cell division in bladder ¹⁹⁷ , breast ¹⁹⁸ , colon ¹⁹⁹ , and other cancers
Cholesterylation 	Hedgehog signaling activation	Facilitate tumorigenesis and cancer growth in prostate ²⁰⁰ , breast ²⁰¹ , bladder ²⁰² , and other cancers
Myristoylation 	Membrane localization Autoinhibition	Carcinogenesis in breast ²⁰³ , lung ²⁰⁴ , and other cancers
Palmitoylation 	Plasma membrane localization Partitioning into lipid rafts Protein maturation/quality control	Promote proliferation and invasion in melanoma ¹⁷⁰ , intestinal ²⁰⁵ , and other cancers
Farnesylation 	Membrane localization Conformational change Protein-protein interaction	Promote cell growth, survival and metastasis in lung ²⁰⁶ , myeloid leukemia ²⁰⁷ , pancreatic ²⁰⁸ , and other cancers
Geranylgeranylation 	Membrane localization Protein-protein interaction	Facilitate cell proliferation and migration in lymphoma ¹⁷⁷ , leukemia ²⁰⁹ , and other cancers




Lipidation

A) Natural Lipidation




Lipidated therapeutics


B) Unnatural Lipidation




Lipidated therapeutic anchors into membrane



Lipidated therapeutic binds to soluble proteins





Lipidated therapeutic binds to membrane-bound proteins




Lipidated therapeutic is more cell permeable

Lipidated therapeutics

Escape from Human Immunodeficiency Virus Type 1 (HIV-1)



chromosome



Fig_11-05 *Genetics, Second Edition* © 2005 W.H. Freeman and Company

Drosophila nucleosome core

H2A, H2B, H3, and H4








Figure 4.38 How Proteins Work (©2012 Garland Science)

Modification



Effects



modification state


"meaning"



heterochromatin formation,
gene silencing



gene expression



gene expression



silencing of Hox genes,
X chromosome inactivation

effects




Figure 4.39 How Proteins Work (©2012 Garland Science)




Figure 4.40 How Proteins Work (©2012 Garland Science)

Protein quality control





Figure 4.12.1 How Proteins Work (©2012 Garland Science)




Figure 4.41 How Proteins Work (©2012 Garland Science)





Figure 4.13.1 How Proteins Work (©2012 Garland Science)

Pancreatic β cells



- In pancreatic **β -cells**, glucose release requires **Ca²⁺ influx**.

- Low** blood glucose:

- $\downarrow$ ATP $\rightarrow$ $\uparrow$ K⁺ efflux $\rightarrow$ $\uparrow$ hyperpolarization $\rightarrow$ $\downarrow$ Ca²⁺ influx $\rightarrow$ $\downarrow$ insulin secretion

- Elevated** blood glucose

- $\uparrow$ ATP $\rightarrow$ $\downarrow$ K⁺ efflux $\rightarrow$

self-splicing polypeptides





Figure 4.10.1 How Proteins Work (©2012 Garland Science)




A biotechnological application of inteins. the fusion of domain 1 with an intein (created genetically by fusing the genes) is induced to self-cleave by addition of an external thiol, RSH.


The thioester thus formed is subsequently cleaved by a second domain starting in a cysteine, to give a two-domain protein.

Figure 4.10.2 How Proteins Work (©2012 Garland Science)


A)



B)





C)




Catalysis

free energy profiles of uncatalyzed and enzyme-catalyzed reactions


(A)



(B)




isomer




D-(+)-glyceraldehyde

Figure 1.1.1 Molecular Biology of Assemblies and Machines (© Garland Science 2016)




L-(-)-glyceraldehyde




D-erythrose

Figure 1.1.2 Molecular Biology of Assemblies and Machines (© Garland Science 2016)



D-threose

enantiomer





thalidomide

Figure 1.1.3 Molecular Biology of Assemblies and Machines (© Garland Science 2016)

diastereoisomer


r

catalytic mechanism of the pancreatic protease chymotrypsin





Substrate specificity

selectivity




Hydrophobic amino acids

Figure 9-1
Biochemistry, Sixth Edition
© 2007 W.H. Freeman and Company



Pocket

F Phenylalanine, Y Tyrosin, W Tryptophan, M Methionine
K Lysine, R Arginine
A Alanine, S Serine



related serine peptidases

Specificity pocket mutations leading to acquisition of tryptic activity

