




Membrane Proteins





Anatomies of two types of biological cells



Evolution of mitochondria








Membrane Lipids Form Bilayers in Water



(E) name of glycerophospholipid	name of X	formula of X	net charge (at pH 7)
phosphatidic acid	-	-H	-1
phosphatidylethanolamine	ethanolamine	-CH ₂ -CH ₂ -NH ₃ ⁺	0
phosphatidylcholine	choline	-CH ₂ -CH ₂ -N ⁺ (CH ₃) ₃	0
phosphatidylserine	serine	-CH ₂ -CH(NH ₃ ⁺)COO ⁻	-1
phosphatidylglycerol	glycerol	-CH ₂ -CH(OH)-CH ₂ -OH	-1
phosphatidylinositol 4,5-bisphosphate	4,5-bisphosphate		-5
cardiolipin	phosphatidylglycerol		-2

(F) name of sphingolipid	name of X	formula of X
ceramide	-	-H
sphingomyelin	phosphocholine	
neutral glycolipids glucosylcerebroside	glucose	
lactosylceramide	di-, tri-, or tetrasaccharide	
ganglioside GM2	complex oligosaccharide	

Molecular dynamics




Composition of the membranes

Composition of Membranes (% by weight)

Membrane	Protein	Lipid	Carbohydrate
Myelin	20	75	5
Red blood cell ghosts	49	43	8
Plasma membranes			
Liver cells	58	42	(5–10) ^a
Ehrlich ascites	67	33	(5–10) ^a
Amoeba <i>Proteus</i>	25	32	15
Mitochondrion			
Outer membrane	52	48	(2–4) ^a
Inner membrane	76	24	(1–2) ^a
Bacteria (Gram-positive)	75	25	(10) ^a

^aCarbohydrate is due to glycosylated proteins and lipids.

Based on data provided in Guidott G [1972] *Arch Intern Med* 129: 194–301.



Curvature model of membrane vesicles




Figure 2.17 Cell Membranes (© Garland Science 2016)





Figure 2.20 Cell Membranes (© Garland Science 2016)



	P	volume	conformation	curvature	lipid
	>1		reverse cone	negative	PE
	$=1$		cylinder	zero	PC
	<1		cone	positive	lyso PC

Lipid diffusion

lateral diffusion



transverse diffusion (flip-flop)



top view



I



II








III






IV



side view



Structure of a DOPC bilayer



Predicting transmembrane helices



(C)

hydrophobicity amino acid values		
Kyte and Doolittle	Eisenberg	Tanford
Ala: 1.800	Ala: 0.620	Ala: 0.620
Arg: -4.500	Arg: -2.530	Arg: -2.530
Asn: -3.500	Asn: -0.780	Asn: -0.780
Asp: -3.500	Asp: -0.900	Asp: -0.090
Cys: 2.500	Cys: 0.290	Cys: 0.290
Gln: -3.500	Gln: -0.850	Gln: -0.850
Glu: -3.500	Glu: -0.740	Glu: -0.740
Gly: -0.400	Gly: 0.480	Gly: 0.480
His: -3.200	His: -0.400	His: -0.400
Ile: 4.500	Ile: 1.380	Ile: 1.380
Leu: 3.800	Leu: 1.060	Leu: 1.530
Lys: -3.900	Lys: -1.500	Lys: -1.500
Met: 1.900	Met: 0.640	Met: 0.640
Phe: 2.800	Phe: 1.190	Phe: 1.190
Pro: -1.600	Pro: 0.120	Pro: 0.120
Ser: -0.800	Ser: -0.180	Ser: -0.180
Thr: -0.700	Thr: -0.050	Thr: -0.050
Trp: -0.900	Trp: 0.810	Trp: 0.810
Tyr: -1.300	Tyr: 0.260	Tyr: 0.260
Val: 4.200	Val: 1.080	Val: 1.800






Figure 3.15 Cell Membranes (© Garland Science 2016)

Hydropathy plot analysis of protein sequences



Topology and orientation of helical integral membrane proteins






Figure 3.4 Cell Membranes (© Garland Science 2016)

Topology diagrams and X-ray structures for various transmembrane proteins



ELIC, pentameric ligand-gated ion channel



cold receptor







Figure 3.11 Cell Membranes (© Garland Science 2016)

Characteristics of transmembrane α helices




Characteristics of transmembrane α helices

**phospholemman
(2JO1)**



**glycophorin
(1AFO)**



Topology diagrams to probe intra- and extracellular orientation





Figure 3.13 Cell Membranes (© Garland Science 2016)

Transmembrane segment variations






Figure 3.18 Cell Membranes (© Garland Science 2016)

Lipid–protein interactions

human glycoporphin A



(A)



1. protein–water interactions
2. protein–protein interactions
3. protein–lipid tail (hydrophobic core) interactions
4. protein–lipid head group (bilayer interface) interactions

Aromatic side chains as membrane anchors

tyrosine, phenylalanine, tryptophan






Figure 3.21 Cell Membranes (© Garland Science 2016)

(A)

KcsA potassium channel



G Gly, P Pro, L Leu, R Arg


(C)

GGPG-L₆RL₁₃-GPGG (R7)

GGPG-L₉RL₁₀-GPGG (R10)
(down)


GGPG-L₉RL₁₀-GPGG (R10)
(up)

GGPG-L₁₂RL₇-GPGG (R13)




● water ● water near peptide ● phosphate ● carbonyl ● arginine ● leucine

Interactions of helices in lipid bilayers



	$\Delta G = G_2 - G_1$ (kcal/mol)	
	in H_2O	in $(CH_2)_n$
hydrophobic	20	0
AB ion pair	2	20 – 30
AB strong H bond	0	10 – 20
AB H bond	-1	5
conformation entropy	-5	-5

Interhelical hydrogen bonds. In bacteriorhodopsin from *Halobacterium salinarum* Tyr185 (orange) of helix 6 and Asp212 (yellow) of helix 7 form an interhelical hydrogen bond (dashed line).








Aromatic–aromatic interactions. In subunit I of aberrant ba3-cytochrome c oxidase from *Thermus thermophilus* Trp110 of helix 4 interacts with Tyr23 and Leu27 of helix 1, although it is partly exposed to the lipid bilayer.

van der Waals contacts. Small residues (orange and yellow) increase the homodimer interface and allow for extensive van der Waals contacts in human glycophorin A. Hydrogens are shown for Gly residues only.

Salt bridges. Lys358 and Asp237 are crucial for membrane insertion of lac permease from *E. coli* and have been suggested to form a salt bridge within the protein's transmembrane region.

Bind to Membranes with the Help of Covalently Linked Lipids





(D)


Src(2-16)	myristate-GS	S K S K P K D P S Q R R R
MARCKS(151-175)		K K K K R F S F K K S F K L S G F S F K K N K K
HIV-1 Gag (2-31)	myristate-GA	R A S V L S G G E L D R W E K I R L R P G G K K Y K L
K-Ras 4B (174-185)		G K K K K K S K T S C-farnesyl

K Lys, R Arg, G Gly, S Ser, D Asp, E Glu, N Asn,
V Val


Lateral Pressure



Lateral Pressure



Forced unfolding of membrane-bound BR



Protein–lipid complexes




Figure 3.25 Cell Membranes (© Garland Science 2016)

Specific lipid effects

Lipids can act as co-factor that facilitate the folding or stabilise the structure of membrane proteins

Diacylglycerol kinase from E. coli, which requires 1,2-dioleoyl-sn-glycero-phosphoglycerol (DOPG) for proper folding.


Cardiolipin, a four-chain lipid binds to the large mitochondrial membrane protein bovine cytochrome c oxidase and is essential for its function. Cardiolipin is explicitly required for association of cytochrome c oxidase subunits IVa and IVb.

Several membrane-protein crystal structures show tightly bound lipid molecules and provide valuable insights into how these specifically interact with membrane proteins


The function of KvAP channel, a voltage-dependent K⁺-channel, depends on certain lipid species. KvAP senses voltage with the aid of Arg-containing structures located at the membrane interface and pointing into the membrane interior

The functional state of KvAP requires POPE or 1-palmitoyl-2-oleoyl-sn-glycero-phosphoglycerol (POPG) and that phosphate groups play a crucial role, as their enzymatic removal disrupts function

Oligomerization and clustering of membrane proteins







Oligomerization and clustering of membrane proteins



200 nm

Membrane active peptides





melittin

a




Peptide	Amino Acid Sequence *												
Melittin	1	2	3	4	5	6	7	8	9	10	11	12	13
	G	I	G	A	V	L	K	V	L	T	T	G	L
	14	15	16	17	18	19	20	21	22	23	24	25	26
	P	A	L	I	S	W	I	K	R	K	R	Q	Q

* Blue: hydrophobic; Orange: hydrophilic; Red: charged

b



Melittin Insertion into Cell Membranes



α -Helical ionophoric peptides




Figure 3.27c Cell Membranes (© Garland Science 2016)







Figure 3.27d Cell Membranes (© Garland Science 2016)




Amphiphilic helices





apolipoprotein A-I
tandem amphiphilic helices



lipoprotein A-I
tandem amphiphilic helices




nanodisc





nanodisc
with bacteriorhodopsin
trimer


antimicrobial peptides (AMPs)




Cyclic peptides self-assemble into β -sheet type nanotubes




Cyclic peptides self-assemble into β -sheet type nanotubes





A



B



Monotopic membrane proteins



teichoic acid biosynthesis protein F





Figure 3.5 Cell Membranes (© Garland Science 2016)

Monotopic membrane proteins


prostaglandin H2 synthase



PGH₂ Synthase is a heme-containing **dioxygenase**, bound to ER membranes.

(A dioxygenase incorporates **O₂** into a substrate).

PGH₂ Synthase exhibits 2 activities: **cyclooxygenase** & **peroxidase**.



Lipid-anchored proteins





Figure 3.6a Cell Membranes (© Garland Science 2016)




Figure 3.6b Cell Membranes (© Garland Science 2016)


Lipid anchor types and function

Table 3.2 Lipid anchor types and function

Common name	Lipid	Linker	Target membrane	Topology	Reversible	Function
Myristic acid	C14:0	N-term G (internal K)	Any	Cytosolic	No	Targeting
Palmitic acid	C16:0	Cys (K, S, T)	Any	Cytosolic	Yes	Sorting, trafficking
Farnesyl	C15	C-term (CaaX, CXC, XXCC)	Plasma	Cytosolic	No	Sorting, protein interaction
Geranylgeranyl	C20	C-term (CaaX, CXC, XXCC)	Plasma	Cytosolic	No	Sorting, protein interaction
GPI (SPI)	Glycerolipid (sphingolipid)	C-term	Plasma	Extracellular	Yes	Cell communication

GPI, glycosylphosphatidylinositol; SPI, sphingolipid inositol; C-term, C-terminal; N-term, N-terminal.

Lateral diffusion is faster for lipid-anchored than for transmembrane proteins



Peripheral membrane proteins as subunits of protein complexes







Figure 3.8 Cell Membranes (© Garland Science 2016)


Peripheral membrane proteins as subunits of protein complexes



Peripheral lipid-binding membrane proteins



PS



PI(4,5)P₂



Figure 3.9b Cell Membranes (© Garland Science 2016)

Peripheral lipid-binding membrane proteins




Figure 3.9a Cell Membranes (© Garland Science 2016)

pleckstrin homology

Action

